

Progress on COUPP60

Hugh Lippincott, Fermilab

for the PICO/COUPP Collaboration
TAUP, Asilomar, CA
September 11, 2013

PICO Collaboration



C. Amole, M. Besnier,
G. Caria, A. Kamaha,
A. Noble, T. Xie



M. Ardid,
M. Bou-Cabo



D. Asner, J. Hall



D. Baxter, C.E. Dahl, M. Jin

NORTHWESTERN
UNIVERSITY

E. Behnke, H. Borsodi,
C. Harnish, O. Harris,
C. Holdeman, I. Levine,
E. Mann, J. Wells

INDIANA UNIVERSITY
SOUTH BEND



P. Bhattacharjee, M. Das,
S. Seth

Fermilab

S.J. Brice, D. Broemmelsiek,
P.S. Cooper, M. Crisler,
W.H. Lippincott, E. Ramberg,
M.K. Ruschman,
A. Sonnenschein



J.I. Collar, R. Neilson,
A.E. Robinson

Université
de Montréal

F. Debris, M. Fines-Neuschild, C.M. Jackson,
M. Lafrenière, M. Laurin, L. Lessard,
J.-P. Martin, M.-C. Piro, A. Plante, O. Scallan,
N. Starinski, V. Zacek



N. Dhungana, J. Farine,
R. Podviyanuk, U. Wichoski



R. Filgas,
S. Pospisil, I. Stekl



D. Maurya, S. Priya

S. Gagnebin, C. Krauss,
D. Marlisov, P. Mitra





I. Lawson,
E. Vázquez Jáuregui



PICO/COUPP fast compression bubble chamber


- Pressure expansion creates superheated fluid, CF_3I
- I for spin-independent
- F for spin-dependent
- Alternatives - e.g. C_3F_8 (see talk on PICO-2L by R. Neilson, DM II)
- Particle interactions nucleate bubbles
- Cameras see bubbles
- Recompress chamber to reset




Why bubble chambers?

- A lot of effort in dark matter experiments goes into discriminating **electronic recoils (gammas)** vs. **nuclear recoils(WIMPs)**

Xenon TPCs -
Charge to light




Argon - Pulse
shape discrimination



Why bubble chambers?


- By choosing superheat parameters appropriately (temperature and pressure), bubble chambers are blind to electronic recoils (10^{-10} or better)



Why bubble chambers?


- By choosing superheat parameters appropriately (temperature and pressure), bubble chambers are blind to electronic recoils (10^{-10} or better)
- To form a bubble requires two things
 - Enough energy
 - Enough energy density - length scale must be comparable to the critical bubble size

Why bubble chambers?



Why bubble chambers?

- Easy to identify multiple scattering events → Neutron backgrounds
- Easy DAQ and analysis chain
 - Cameras
 - Piezos
- No PMTs, no cryogenics




Why not bubble chambers?

- Threshold detectors - no energy resolution
 - Harder to distinguish some backgrounds, less information about any potential signal
 - Alphas (several MeV) were a big concern
 - Energy threshold calibrations are hard and important
- Bubble chambers are slow - about 30 s of deadtime for every event
 - Overall rate must be low




About those alphas

- Discovery of acoustic discrimination against alphas by PICASSO (Aubin et al, New J. Phys 10:103017, 2008)
- Alphas deposit energy over tens of microns
- Nuclear recoils deposit theirs in tens of nanometers
- In COUPP bubble chambers, alphas are several times louder



The COUPP/PICO program

- COUPP4: A 2-liter chamber run at SNOLAB in 2010-2012
- PICO-2L: see talk from R. Neilson, Session DM 2
- COUPP60: Up to 40 liters, running at SNOLAB now
- PICO-250L (formerly COUPP500): ton-scale G2 experiment




COUPP-4



COUPP4: Acoustic discrimination

- Better than 99.3% rejection against alphas at 16 keV threshold
- Limited by statistics, and backgrounds





COUPP4: Results and sensitivity



- 20 WIMP candidates (6 at 8 keV, 6 at 11 keV, 8 at 16 keV)
- 3 multiple bubble events imply some contribution from neutrons
- U,Th in the piezo-acoustic sensors and the viewports
- Remaining excess of singles at low threshold
- Time clustering
- Correlated with activity at water-CF₃I interface

COUPP4: Results and sensitivity

- Results in PRD 86:052001 (2012)
- Second run in 2012 observed same phenomena (Eric Vazquez Jauregui, Moriond, 2013)




COUPP60

- Engineering run at shallow site in 2010
- Low backgrounds and acoustic discrimination



COUPP60

- Engineering run at shallow site in 2010
- Low backgrounds and acoustic discrimination
- Fluid darkening due to photodissociation of iodine
- Excessive surface rate



COUPP60

- Engineering run at shallow site in 2010
- Low backgrounds and acoustic discrimination
- Fluid darkening due to photodissociation of iodine
- Excessive surface rate
- Solutions tested in second run November, 2011
- Started moving to SNOLAB last summer






Tuesday, September 10, 2013



Tuesday, September 10, 2013




Tuesday, September 10, 2013




Tuesday, September 10, 2013

COUPP60



- Filled with 36.8 kg of CF_3I at end of April
- First bubble observed on May 1 (radon decay)
- Physics data started June 13


COUPP60




- Collected 1378 kg-days of dark matter search data between 10 and 20 keV threshold
- Good live fraction $> 80\%$, no darkening
- > 1500 neutron source events from calibration runs

COUPP60


- August 9, we discovered a small hydraulic leak
- We stopped cycling the chamber, leak closed up
- Most likely scenario is pressure relief opening on compression
- Operations on hold until we can get into the water tank and fix the problem
- Will be running again in a couple of weeks



COUPP60 - the data




- Position reconstruction working well
- Clear set of events on surface and hemisphere
- Not a background, and rate is under control




COUPP60 - the data

Preliminary




- Acoustics are working well
- Calibration data show a narrower distribution than COUPP4
- Alpha discrimination still very strong




COUPP60 - the data

- Analysis still under development
- Good news:
- Bad news:




COUPP60 - the data

- Analysis still under development
- Good news: Zero multiple bubbles, no neutrons. Limit on neutron rate is factor 3 below observed rate in COUPP4
- Bad news:




COUPP60 - the data

- Analysis still under development
- Good news: Zero multiple bubbles, no neutrons. Limit on neutron rate is factor 3 below observed rate in COUPP4
- Bad news: Population of events that sound like nuclear recoils but are clearly not WIMPs
- Silver lining:




COUPP60 - the data

- Analysis still under development
- Good news: Zero multiple bubbles, no neutrons. Limit on neutron rate is factor 3 below observed rate in COUPP4
- Bad news: Population of events that sound like nuclear recoils but are clearly not WIMPs
 - Silver lining: statistics - we can actually study them in detail
 - Early indications confirm a slightly different acoustic distribution and similar timing and spatial correlations to COUPP4 background for at least some fraction of events





COUPP60 - future plans

- Now in Phase I: Continue running until end of year at different thresholds, fully characterize the detector, first physics result
- Phase II: Upgrade to full 75 kg target with second set of cameras, collect 10000 kg-days of data
- Planned three year run with potential target swap as new information comes in



Conclusion

- COUPP60 operating well with a good live fraction (at least until the leak)
- Acoustics continue to provide good discrimination power in the larger chamber
- Still some work to do to understand the detector and extract physics - but it's early yet and who wanted it to be easy?



Extra

Detour: Threshold and efficiency



- Threshold determined from Seitz, Phys. of Fluids **1**, 2 (1958)

$$p_v - p_l = \frac{2\sigma}{r_c}$$
$$E_{th} = \underbrace{4\pi r_c^2 \left(\sigma - T \frac{\partial \sigma}{\partial T} \right)}_{\text{Surface energy}} + \underbrace{\frac{4}{3}\pi r_c^3 \rho_v h}_{\text{Latent heat}}$$

- Energy deposition E_{th} within length R_c will nucleate a bubble (Hot Spike model)
- Theory assumes a step function above threshold

Detour: Threshold and efficiency



$$\text{Rate} = \int \text{WIMP recoil spectrum} \times \text{Bubble nucleation efficiency}$$



- Effect of threshold shape depends on target, WIMP mass

Detour: Threshold and efficiency



$$\text{Rate} = \int \text{WIMP recoil spectrum} \times \text{Bubble nucleation efficiency}$$



- Effect of threshold shape depends on target, WIMP mass

Detour: Threshold and efficiency

$$\text{Rate} = \int \text{WIMP recoil spectrum} \times \text{Bubble nucleation efficiency}$$



- Effect of threshold shape depends on target, WIMP mass

Detour: Threshold and efficiency



- Threshold determined from Seitz, Phys. of Fluids **1**, 2 (1958)

$$p_v - p_l = \frac{2\sigma}{r_c}$$
$$E_{th} = \underbrace{4\pi r_c^2 \left(\sigma - T \frac{\partial \sigma}{\partial T} \right)}_{\text{Surface energy}} + \underbrace{\frac{4}{3}\pi r_c^3 \rho_v h}_{\text{Latent heat}}$$

- Energy deposition E_{th} within length r_c will nucleate a bubble (Hot Spike model)
- Theory assumes a step function above threshold
- Needs calibration



Detour: Threshold and efficiency

- Complicated by molecule, CF_3I
- Recall that the recoil track length L must be comparable to the bubble radius R_c




Carbon and fluorine

- Use neutron calibration sources at SNOLAB
- Compare MCNP-predicted rates of single, double, triple and quadruple bubble events with observation
- Data show a shortfall of events compared to simulation of the Seitz Model- i.e. the threshold is not a step function



What about iodine?


- Main sensitivity to spin independent dark matter from iodine
 - 85% of neutron source interactions are with C and F
- Heavy radon daughter nuclei are a proxy and are step-like



- We really need a direct calibration

COUPP Iodine Recoil Threshold Experiment


- Bubble chambers are insensitive to MIPs
- Elastic scattering of charged particles can be tracked with very high precision



$$T = E_{\text{recoil}} = \frac{(p\theta)^2}{2m_r}$$

COUPP Iodine Recoil Threshold Experiment


- Provides event by event energy information bubble chambers normally can't provide
- 75% of elastic scattering events with 12 GeV pions at energies relevant to dark matter involve iodine



$$T = E_{\text{recoil}} = \frac{(p\theta)^2}{2m_r}$$

COUPP Iodine Recoil Threshold Experiment

- Test beam at Fermilab with a silicon pixel telescope
- Designed a new test tube sized bubble chamber




Hydraulics

Piezo-acoustic sensor




Beam tube

Water bath



COUPP Iodine Recoil Threshold Experiment

- Beam run at Fermilab in March, 2012



COUPP Iodine Recoil Threshold Experiment

- Analysis shows that iodine threshold is very close to a step function at the predicted energy
 - Limited by resolution (MCS) and statistics

